

***Lactobacillus paracasei* Shirota: studi chiave**

***Lactobacillus paracasei* Shirota (L. casei Shirota, LcS)** è l'esclusivo ceppo probiotico contenuto nelle tre referenze Yakult: Yakult Original, Yakult Light e Yakult Plus.

Per maggiori informazioni, è possibile visitare il sito www.scienceforhealth.it, riservato a medici e professionisti della salute.

Per contattare il **Dipartimento Scientifico di Yakult Italia** è possibile scrivere a science@yakult.it oppure telefonare al numero 02831281.

Indice

1. Sopravvivenza nel tratto gastro-intestinale
2. Funzioni gastrointestinali (stipsi, IBS, IBD, NEC, epatopatie, intolleranza al lattosio, H. pylori)
3. Diarrea infettiva, diarrea associata alla terapia antibiotica (AAD) e diarrea associata a *Clostridium difficile* (CDAD)
4. Altre infezioni
5. Malattie metaboliche e rischio cardiovascolare (diabete mellito, resistenza insulinica, ipertensione arteriosa)
6. Asse intestino-cervello
7. Cancro (colon-retto, vescica, polmone, mammella)
8. Sistema immunitario
9. Miscellanea (nutrizione enterale, sport, ...)

1. Sopravvivenza nel tratto gastro-intestinale

Dynamic analysis of human small intestinal microbiota after an ingestion of fermented milk by small-intestinal fluid perfusion using an endoscopic retrograde bowel insertion technique. Takada T, Chinda D, Mikami T, et al. *Gut Microbes*. 2020;11(6):1662-1676. doi:10.1080/19490976.2020.1766942

[\[Free paper\]](#) [\[PubMed\]](#)

Recovery of *Lactobacillus casei* strain Shirota (LcS) from the intestine of healthy Vietnamese adults after intake of fermented milk. Mai TT, Hop DV, Anh TT et al. *Asia Pacific Journal of Clinical Nutrition* 2017;26(1):72-77

[\[Free paper\]](#) [\[PubMed\]](#)

Survival of *Lactobacillus casei* strain Shirota in the intestines of healthy Chinese adults. Wang R, Chen S, Jin J et al. *Microbiology Immunology* 2015;59(5):268-76

[\[PubMed\]](#)

Intestinal Microbiota Profiles of Healthy Pre-School and School-Age Children and Effects of Probiotic Supplementation. Wang et al. *Ann Nutr Metab* 2015;67(4): 257-66.

[\[PubMed\]](#)

Survivability of a probiotic *Lactobacillus casei* in the gastrointestinal tract of healthy human volunteers and its impact on the faecal microflora. Tuohy KM et al. *J Appl Microbiol* 2007;102(4): 1026-1032.

[\[PubMed\]](#)

Survival of a probiotic, *Lactobacillus casei* strain Shirota in the gastrointestinal tract: selective isolation from faeces and identification using monoclonal antibodies. Yuki N, Watanabe K, Mike A et al. *International Journal of Food Microbiology* 1999;48:51-57.

[\[PubMed\]](#)

The effect of consumption of milk fermented by *Lactobacillus casei* strain Shirota on the intestinal immune parameters in humans. *Eur J Clin Nutr*. 1998 Dec;52(12):899-907

[\[PubMed\]](#)

The probiotic *Lactobacillus paracasei* strain Shirota (LcS) in a fermented milk beverage survives the gastrointestinal tract of generally healthy U.S. Adults. Cook CM, Makino H, Kato K, Blonquist T, Derrig L, Shibata H. *Int J Food Sci Nutr*. 2023 Sep;74(5):645-653. doi: 10.1080/09637486.2023.2246693. Epub 2023 Aug 16. PMID: 37584253.

[\[PubMed\]](#)

2. Funzioni gastrointestinali (stipsi, IBS, IBD, NEC, epatopatie, intolleranza al lattosio, H. pylori)

Probiotic *Lactobacillus casei* Shirota prevents acute liver injury by reshaping the gut microbiota to alleviate excessive inflammation and metabolic disorders. Yan R, Wang K, Wang Q, Jiang H, Lu Y, Chen X, Zhang H, Su X, Du Y, Chen L, Li L, Lv L. *Microb Biotechnol.* **2021** Jan 25. doi: 10.1111/1751-7915.13750.

[\[PubMed\]](#)

Effect of *Lactobacillus casei* on lipid metabolism and intestinal microflora in patients with alcoholic liver injury. Li X, Liu Y, Guo X, Ma Y, Zhang H, Liang H. *Eur J Clin Nutr.* **2021** Jan 29. doi: 10.1038/s41430-020-00852-8

[\[PubMed\]](#)

The efficacy and safety of probiotics for patients with constipation-predominant irritable bowel syndrome: A systematic review and meta-analysis based on seventeen randomized controlled trials. Wen Y, Li J, Long Q, Yue CC, He B, Tang XG. *Int J Surg.* **2020**;79:111-119. doi:10.1016/j.ijisu.2020.04.063

[\[PubMed\]](#)

Are probiotics useful in the treatment of chronic idiopathic constipation in adults? A review of existing systematic reviews, meta-analyses, and recommendations. Kamiński M, Skonieczna-Żydecka K, Łoniewski I, Koulaouzidis A, Marlicz W. *Prz Gastroenterol.* **2020**;15(2):103-118. doi:10.5114/pg.2019.86747

[\[PubMed\]](#)

A Double-Blind, Randomized Placebo-Controlled Trial of Probiotic *Lactobacillus casei* Shirota in Stable Cirrhotic Patients. Macnaughtan J, Figorilli F, García-López E, et al. *Nutrients.* **2020**;12(6):1651. Published 2020 Jun 2. doi:10.3390/nu12061651

[\[Free paper\]](#) [\[PubMed\]](#)

Modulatory Effects of Gut Microbiota on Constipation: The Commercial Beverage Yakult Shapes Stool Consistency. Min Chen, Xujun Ye, Dan Shen, Chunwei Ma. *J Neurogastroenterol Motil.* **2019** Jul; 25(3): 475–477.

[\[PubMed\]](#)

***Lactobacillus casei* Strain Shirota alleviates constipation in adults by increasing the pipercolinic acid level in the gut.** Yangwenshan Ou, Shanbin Chen, Fazheng Ren, Ming Zhang, Shaoyang Ge, Huiyuan Guo, Hao Zhang and Liang Zhao. *Front Microbiol* **2019**; 10: 324.

[\[PubMed\]](#)

Differential Effects of *Lactobacillus casei* Strain Shirota on Patients With Constipation Regarding Stool Consistency in China. Shanbin Chen, Yangwenshan Ou, Liang Zhao, Yang Li, Zhenxing Qiao, Yanling Hao, Fazheng Ren. *J Neurogastroenterol Motil.* **2019**;25:1

[\[PubMed\]](#)

Fermented milk containing *Lactobacillus casei* strain Shirota preserves the diversity of the gut microbiota and relieves abdominal dysfunction in healthy medical students exposed to

academic stress. Kato-Katoaka A, Nishida K, Takada M *et al. Applied and Environmental Microbiology* **2016**;82(12):3649-58.

[\[PubMed\]](#)

The effectiveness of *Lactobacillus* beverage in controlling infections among the residents of an aged care facility: a randomized placebo-controlled double-blind trial.

Nagata S, Asahara T, Wang C *et al. Annals of Nutrition & Metabolism* **2016**;68(1):51-59.

[\[PubMed\]](#)

Efficacy of *Lactobacillus casei* Shirota for patients with irritable bowel syndrome. Thijssen AY, Clemens CH, Vankerckhoven V *et al. European Journal of Gastroenterology & Hepatology* **2015**;28(1):8-14.

[\[PubMed\]](#)

Probiotic *Lactobacillus casei* Shirota improves kidney function, inflammation and bowel movements in hospitalized patients with acute gastroenteritis – A prospective study. Akoglu B, Loytved A, Nuiding H, *et al. Journal of Functional Foods* **2015**;17:305-313.

[\[Abstract\]](#)

Effect of fermented milk containing *Lactobacillus casei* strain Shirota on constipation-related symptoms and haemorrhoids in women during puerperium. Sakai T, Kubota H, Gawad A *et al. Beneficial Microbes* **2015**;6(3):253-262.

[\[PubMed\]](#)

Improving the bowel habits of elderly residents in a nursing home using probiotic fermented milk. Van den Nieuwboer M, Klomp-Hogeterp A, Verdoorn S *et al. Beneficial Microbes* **2015**;6(4):397-403.

[\[Free paper\]](#) [\[PubMed\]](#)

Maintenance of healthy intestinal microbiota in women who regularly consume probiotics. Tsuji H, Chonan O, Suyama Y *et al. Int J Probiotics Prebiotics* **2014**;9(1/2):31-38.

[\[Abstract\]](#)

A probiotic fermented milk drink containing *Lactobacillus casei* strain Shirota improves stool consistency of subjects with hard stools. Tilley L, Keppens K, Kushiro A *et al. Int J Probiotics Prebiotics* **2014**;9(1/2):23-30.

[\[Available on request\]](#)

Beneficial effects of long-term consumption of a probiotic combination of *Lactobacillus casei* Shirota and *Bifidobacterium breve* Yakult may persist after suspension of therapy in lactose-intolerant patients. Almeida CC, Lorena SLS, Pavan CR *et al. Nutrition in Clinical Practice* **2012**;27(2):247-251.

[\[PubMed\]](#)

Successful treatment of primary sclerosing cholangitis with a steroid and a probiotic. Shimizu M, Iwasaki H, Mase S *et al. Case Reports in Gastroenterology* **2012**;6:249-253.

[\[Free paper\]](#) [\[PubMed\]](#)

Efficacy of *Bifidobacterium breve* and *Lactobacillus casei* oral supplementation on necrotizing enterocolitis in very-low-birth-weight preterm infants: a double-blind, randomized, controlled trial. Braga TD, da Silva GA, de Lira PIC *et al. American Journal of Clinical Nutrition* **2011**;93:81-86.

[\[Free paper\]](#) [\[PubMed\]](#)

Efficacy of *Lactobacillus casei* treatment on small bowel injury in chronic low-dose aspirin users: a pilot randomized controlled study. Endo H, Higurashi T, Hoson K *et al.* *Journal of Gastroenterology* **2011**;46:894-905.

[\[PubMed\]](#)

Effect of *Lactobacillus casei* Shirota on colonic transit time in patients with chronic constipation. Krammer H.-J, von Seggern H, Schaumburg J *et al.* *Coloproctology* **2011**;33:109-113.

[\[Article\]](#)

Fermented milk containing *Lactobacillus casei* strain Shirota reduces incidence of hard or lumpy stools in healthy population. Sakai T, Makino H, Ishikawa E *et al.* *International Journal of Food Sciences & Nutrition* **2011**;62(4):23-30.

[\[PubMed\]](#)

Effects of a probiotic fermented milk beverage containing *Lactobacillus casei* strain Shirota on defecation frequency, intestinal microbiota, and the intestinal environment of healthy individuals with soft stools. Matsumoto K, Takada T, Shimizu K *et al.* *Journal of Bioscience & Bioengineering* **2010**;110(5):547-52.

[\[PubMed\]](#)

Probiotic effects on intestinal fermentation patterns in patients with irritable bowel syndrome. Barrett JS, Canale KEK, Geary RB *et al.* *World Journal of Gastroenterology* **2008**;14(32):5020-5024.

[\[PubMed\]](#)

Beneficial effects of *Lactobacillus casei* in ulcerative colitis: a pilot study. Mitsuyama K, Matsumoto S, Yamasaki H *et al.* *Journal of Clinical Biochemistry and Nutrition* **2008**;43(Suppl 1):78-81.

Effect of probiotic treatment on deranged neutrophil function and cytokine responses in patients with compensated alcoholic cirrhosis. Stadlbauer V, Mookerjee RP, Hodges S *et al.* *Journal of Hepatology* **2008**;48:945-951.

[\[PubMed\]](#)

High dose probiotic and prebiotic co-therapy for remission induction of active Crohn's disease. Fujimori S, Tatsuguchi A, Gudis K *et al.* *Journal of Gastroenterology & Hepatology* **2007**;22: 1199-1204.

[\[PubMed\]](#)

The effect of a probiotic milk product containing *Lactobacillus casei* strain Shirota on the defecation frequency and the intestinal microflora of sub-optimal health state volunteers: a randomized placebo-controlled cross-over study. Matsumoto K, Takada T, Shimizu K *et al.* *Bioscience Microflora* **2006**;25(2):39-48.

[\[Available on request\]](#)

The effects of a synbiotic fermented milk beverage containing *Lactobacillus casei* strain Shirota and transgalactosylated oligosaccharides on defecation frequency, intestinal microflora, organic acid concentrations, and putrefactive metabolites of sub-optimal health state volunteers: a randomized placebo-controlled cross-over study. Shioiri T, Yahagi K, Nakayama S *et al.* *Bioscience Microflora* **2006**;25(4):137-146.

[\[Free paper\]](#)

Effect of frequent consumption of a *Lactobacillus casei*-containing milk drink in *Helicobacter pylori*-colonized subjects. Cats A, Kuipers EJ, Bosscherts MAR *et al. Alimentary Pharmacology & Therapeutics* **2003**;17(3):429-435.

[\[Free paper\]](#)

Probiotic beverage containing *Lactobacillus casei* Shirota improves gastrointestinal symptoms in patients with chronic constipation. Koebnick C, Wagner I, Leitsmann P *et al. Canadian Journal of Gastroenterology* **2003**;17(11):655 – 659.

[\[Available on request\]](#) [\[Free paper\]](#) [\[PubMed\]](#)

Effect of administration of *Lactobacillus casei* Shirota on sodium balance in an infant with short bowel syndrome. Candy DCA, Densham L, Lamont LS *et al. Journal of Pediatric Gastroenterology & Nutrition* **2001**;32(4):506-508.

[\[PubMed\]](#)

3. Diarrea infettiva, diarrea associata alla terapia antibiotica (AAD) e diarrea associata a *Clostridium difficile* (CDAD)

A study into the effect of *Lactobacillus casei* Shirota in preventing antibiotic associated diarrhoea including *Clostridioides difficile* infection in patients with spinal cord injuries: a multicentre randomised, double-blind, placebo-controlled trial.

Wong S *et al., EClinicalMedicine* **2021**; Sep 11;40:101098

[\[PubMed\]](#)

Use of probiotics in preventing antibiotic-associated diarrhoea and *Clostridium difficile* associated diarrhoea in spinal injury centres: an international multicentre study. Wong S, Saif M, O'Driscoll J *et al. International Journal of Probiotics Prebiotics* **2015**;10(23):85-90.

[\[Free paper\]](#)

Prospective randomized double-blind study of probiotic milk drink in reducing the incidence of antibiotic-associated diarrhoea and *Clostridium difficile* diarrhoea.

A. Helps, E. Bell and R. Mactier. *International Journal of Probiotics Prebiotics* **2015**;10(4):145-152

[\[Available on request\]](#)

Probiotic treatment for the prevention of antibiotic-associated diarrhoea in geriatric patients: A multicentre randomised controlled pilot study.

Wright K, Wright H, Murray M. *Australasian Journal Ageing*. **2015** Mar;34(1):38-42

[\[PubMed\]](#)

A *Lactobacillus casei* Shirota probiotic drink reduces antibiotic-associated diarrhoea in patients with spinal cord injuries: a randomized controlled trial. Wong S, Jamous A, O'Driscoll J *et al. British Journal of Nutrition* **2014**;111(4):672-678.

[\[Available on request\]](#)

Prevention of relapse following *Clostridium difficile* infection using probiotic *Lactobacillus casei* Shirota. Lee LYW, Golmohamad R, MacFaul G. *International Journal of Probiotics & Prebiotics* **2013**;8(4):145-148.

[\[Available on request\]](#)

Effects of antibiotic therapy on the gastrointestinal microbiota and the influence of *Lactobacillus casei*. Pirker A, Stockenhuber A, Remely M *et al* () *Food & Agricultural Immunology* **2013**;24(3):315-330.

[\[Available on request\]](#) [\[Abstract\]](#)

Effects of the continuous intake of *Lactobacillus casei* strain Shirota-fermented milk on risk management of long-term inpatients at health service facilities for the elderly. Bian L, Nagata S, Asahara T *et al*. *International Journal of Probiotics and Prebiotics* **2011**;6(2):123-132.

[\[Abstract\]](#)

Effect of the continuous intake of probiotic-fermented milk containing *Lactobacillus casei* strain Shirota on fever in a mass outbreak of norovirus gastroenteritis and the faecal microflora in a health service facility for the aged. Nagata S, Asahara T, Ohta T *et al*. *British Journal of Nutrition* **2011**;106(4):549-56

[\[PubMed\]](#)

Effects of perioperative synbiotic treatment on infectious complications, intestinal integrity and faecal flora and organic acids in hepatic surgery with or without cirrhosis. Usami M, Miyishi M, Kanbara Y *et al*. *Journal of Parenteral and Enteral Nutrition* **2011**;15:317-328.

[\[Free paper\]](#) [\[PubMed\]](#)

Role of probiotic in preventing acute diarrhoea in children: a community-based, randomized, double-blind placebo-controlled field trial in an urban slum. Sur D, Manna B, Niyogi SK, Ramamurthy T *et al*. *Epidemiology and Infection* **2010**;139(6):919-926.

[\[Available on request\]](#) [\[PubMed\]](#)

The potential of probiotic fermented milk products in reducing risk of antibiotic-associated diarrhoea and *Clostridium difficile* disease. Lewis JN, Thomas LV & Weir W. *International Journal of Dairy Technology* **2009**;62(4):461-471.

[\[Free paper\]](#)

Preventing antibiotic associated diarrhoea using a probiotic *Lactobacillus Casei* preparation

Kamhuber C, Hörst-Kollmann S, Rek A, Eckhardt G, Zauner A, Gabriel P, Prager E, Mach K, Stockenhuber F. *Z Gastroenterol.* **2009**; 47 - P1

[\[Abstract\]](#)

Resolution of cryptosporidiosis with probiotic treatment. Pickerd N & Tuthill D. *Postgraduate Medical Journal* **2004**;80:112-113.

[\[Free paper\]](#) [\[PubMed\]](#)

Clinical effect of biolactis powder, a *Lactobacillus casei* preparation, on rotavirus-induced pediatric enteritis. Sugita T & Togawa M. *Japanese Journal of Pediatrics* 1994;47:2755-2762.

[\[Abstract\]](#)

4. Altre infezioni

Synbiotics modulate gut microbiota and reduce enteritis and ventilator-associated pneumonia in patients with sepsis: a randomized controlled trial. Shimizu K, Yamada T, Ogura H, Mohri T, Kiguchi T, Fujimi S, Asahara T, Yamada T, Ojima M, Ikeda M, Shimazu T. *Crit Care*. 2018 Sep 27;22(1):239. doi: 10.1186/s13054-018-2167-x

[\[Free paper\]](#)

Daily fermented milk with *Lactobacillus casei* strain Shirota reduces the incidence and duration of upper respiratory tract infections in healthy middle-aged office workers. Shida K, Sato T, Iizuka R *et al.* *European Journal of Nutrition* 2017;56(1): 45-53

[\[Free paper\]](#) [\[PubMed\]](#)

The effectiveness of *Lactobacillus* beverage in controlling infections among the residents of an aged care facility: a randomized placebo-controlled double-blind trial. Nagata S, Asahara T, Wang C, Suyama Y *et al.* *Annals of Nutrition & Metabolism* 2016;68(1):51-59.

[\[PubMed\]](#)

Randomized study of the effect of synbiotics during neoadjuvant chemotherapy on adverse events in esophageal cancer patients. Motoori M, Yano M, Miyata H *et al.* *Clinical Nutrition* 2015;36(1):93-99

[\[PubMed\]](#)

Effect of probiotic supplement on cytokine levels in HIV-infected individuals: a preliminary study. Falasca K, Vecchiet J, Ucciferri *et al.* *Nutrients* 2015;7:8335-8347.

[\[Free paper\]](#) [\[PubMed\]](#)

Decreased duration of acute upper respiratory tract infections with daily intake of fermented milk: A multicentre, double-blinded, randomized comparative study in users of day care facilities for the elderly population. Fujita R, Iimuro S, Shinozaki T *et al.* *American Journal of Infection Control* 2013;41(12):1231-1235.

[\[PubMed\]](#)

Yakult: a role in combating multi-drug resistant *Pseudomonas aeruginosa*? Thomson CH, Hassan I, Dunn K. *Journal of Wound Care* 2012;21(11):566-569.

[\[Abstract\]](#)

Perioperative synbiotic treatment to prevent infectious complications in patients after elective living donor liver transplantation. A prospective randomised study. Eguchi S, Takatsuki M, Hidaka M *et al.* *American Journal of Surgery* 2010;201(4):498-502.

[\[PubMed\]](#)

Daily probiotic's (*Lactobacillus casei* Shirota) reduction of infection incidence in athletes. Gleeson M, Bishop NC, Oliveira M, Tauler P. *International Journal of Sport Nutrition & Exercise Metabolism* **2011**;21:55-64.

[\[Available on request\]](#) [\[PubMed\]](#)

Perioperative synbiotic treatment to prevent postoperative infectious complications in biliary cancer surgery: A randomized controlled trial. Sugawara G *et al. Annals of Surgery* **2006**;244(5):706-714.

[\[Free paper\]](#)

Synbiotics reduce postoperative infectious complications: a randomized controlled trial in biliary cancer patients undergoing hepatectomy. Kanazawa H, Nagino M, Nishio H *et al. Langenbeck's Arch Surgery* **2005**;390(2): 104-113.

[\[PubMed\]](#)

In vitro and in vivo inhibition of *Helicobacter pylori* by *Lactobacillus casei* strain Shirota. Sgouras D, Maragkoudakis P, Petraki K *et al. Applied & Environmental Microbiology* **2004**;70(1):518-26.

[\[Free paper\]](#) [\[PubMed\]](#)

A prospective uncontrolled trial of fermented milk drink containing viable *Lactobacillus casei* strain Shirota in the treatment of HTLV-1 associated myelopathy / tropical spastic paraparesis. Matsuzaki T, Saito M, Usuku K *et al. Journal of the Neurological Sciences* **2005**;273:75-81.

[\[Abstract\]](#)

5. Malattie metaboliche e rischio cardiovascolare (diabete mellito, resistenza insulinica, ipertensione arteriosa)

Effects of Synbiotic Supplementation on Chronic Inflammation and the Gut Microbiota in Obese Patients with Type 2 Diabetes Mellitus: A Randomized Controlled Study. Kanazawa A, Aida M, Yoshida Y, Kaga H, Katahira T, Suzuki L, Tamaki S, Sato J, Goto H, Azuma K, Shimizu T, Takahashi T, Yamashiro Y, Watada H. *Nutrients*. **2021** Feb 8;13(2):558.

[\[Free paper\]](#) [\[PubMed\]](#)

Effect of probiotic and synbiotic formulations on anthropometrics and adiponectin in overweight and obese participants: A systematic review and meta-analysis of randomized controlled trials. Cao S, Ryan P.M, Salehisahlabadi A. *Journal of King Saud University – Science* Volume 32, Issue 2, March **2020**, Pages 1738-1748

[\[Free paper\]](#)

Effect of *Lactobacillus casei* strain Shirota-fermented milk on metabolic abnormalities in obese prediabetic Japanese men: a randomised, double-blind, placebo-controlled trial. Naito E, Yoshida, Kunihiro S *et al. Bioscience of Microbiota, Food and Health* **2018**;Vol. 37 (1), 9–18, 2018

[\[PubMed\]](#)

Habitual intake of fermented milk products containing *Lactobacillus casei* strain Shirota and a reduced risk of hypertension in older people. Aoyagi Y, Park S, Matsubara S *et al. Beneficial Microbes* **2016**;8(1): 23-29.

[\[Free paper\]](#)

Probiotic supplementation prevents high-fat, overfeeding-induced insulin resistance in human subjects. Hulston CJ, Churnside AA, Venables MC. *British Journal of Nutrition* **2015**;113(4):596-602

[\[Free paper\]](#) [\[PubMed\]](#)

Antihypertensive effect of extracts of *Lactobacillus casei* in patients with hypertension. Nakajima K, Hata Y, Osono Y *et al. Journal of Clinical Biochemistry & Nutrition* **1995**;(18):181-187.

[\[Free paper\]](#) [\[Abstract\]](#)

6. Asse intestino-cervello

Effect of *Lacticaseibacillus paracasei* strain Shirota supplementation on clinical responses and gut microbiome in Parkinson's disease. Yang X, He X, Xu S, Zhang Y, Mo C, Lai Y, Song Y, Yan Z, Ai P, Qian Y, Xiao Q. *Food Funct.* **2023** Jul 31;14(15):6828-6839. doi: 10.1039/d3fo00728f. PMID: 37470081 [\[PubMed\]](#)

Effects of Probiotics on Anxiety, Stress, Mood and Fitness of Badminton Players. Salleh RM, Kuan G, Aziz MNA, Rahim MRA, Rahayu T, Sulaiman S, Kusuma DWY, Adikari AMGCP, Razam MSM, Radhakrishnan AK, Appukutty M. *Nutrients.* **2021** May 24;13(6):1783.

[\[Free paper\]](#) [\[PubMed\]](#)

Effect of *Lacticaseibacillus paracasei* Strain Shirota on Improvement in Depressive Symptoms, and Its Association with Abundance of Actinobacteria in Gut Microbiota.

Otaka M *et al., Microorganisms* **2021**, 9(5):1026.

[\[Free paper\]](#)

Correlation Between Lactic Acid Bacteria Beverage Intake and Stress Resilience. Kato Y, Shimomura Y, Takada Y, Furuyashiki T. *Kobe J. Med. Sci.*, **2021** Vol. 67, No. 1, pp. E1-E6, 202

[\[Free paper\]](#)

Effects of Fermented Milk Containing *Lacticaseibacillus paracasei* Strain Shirota on Constipation in Patients with Depression: A Randomized, Double-Blind, Placebo-Controlled Trial. Zhang X, Chen S, Zhang M, Ren F, Ren Y, Li Y, Liu N, Zhang Y, Zhang Q, Wang R. *Nutrients.* **2021** Jun 29;13(7):2238.

[\[Free paper\]](#) [\[PubMed\]](#)

Beneficial effects of *Lactobacillus casei* strain Shirota on academic stress-induced sleep disturbance in healthy adults: a double-blind, randomised, placebo-controlled trial. Takada M, Nishida K, Gondo Y *et al. Beneficial Microbes* **2017**;8(2):153-162

[\[Free paper\]](#) [\[PubMed\]](#)

Probiotic *Lactobacillus casei* strain Shirota relieves stress-associated symptoms by modulating the gut–brain interaction in human and animal models. Takada M, Nishida K, Kataoka-Kato A *et al. Neurogastroenterology & Motility* **2016**;28(7):1027-1036

[\[PubMed\]](#)

Fermented milk containing *Lactobacillus casei* strain Shirota preserves the diversity of the gut microbiota and relieves abdominal dysfunction in healthy medical students exposed to academic stress. Kato-Kataoka, A. *et al. Appl. Environ. Microbiol.* **2016**;82, 3649–3658 [\[Free paper\]](#)

Use of probiotics for the treatment of constipation in Parkinson’s disease patients. Cassani E, Privitera G, Pezzoli G *et al. Minerva Gastroenterologica e Dietologica* **2011**;57:117-121.

[\[PubMed\]](#)

A randomized, double-blind, placebo-controlled pilot study of a probiotic in emotional symptoms of chronic fatigue syndrome. Rao AV, Bested AC, Beaulne TM *et al. Gut Pathogens* **2009**;1:6 doi:10.1186/1757-4749-1-6

[\[Free paper\]](#) [\[PubMed\]](#)

Impact of consuming a milk drink containing a probiotic on mood and cognition. Benton D, Williams C, Brown A. *European Journal of Clinical Nutrition*, **2007**;61:355–361.

[\[PubMed\]](#)

7. Cancro (colon-retto, vescica, polmone, mammella)

Probiotic beverage with soy isoflavone consumption for breast cancer prevention: a case control study. Toi M, Hirota S, Tomotaki A, *et al. Current Nutrition & Food Science* **2013**;9(4):194-200.

[\[Available on request\]](#) [\[Free paper\]](#)

Probiotics enhance the clearance of human papillomavirus-related lesions: a prospective controlled pilot study. Verhoeven V, Renard N, Makar A *et al. European Journal of Cancer* **2012**;22(1):46-51.

[\[Available on request\]](#) [\[PubMed\]](#)

Prevention of recurrence with Epirubicin and *Lactobacillus casei* after transurethral resection of bladder cancer. Naito S, Koga H, Yamaguchi A *et al. Journal of Urology* **2008**;179:485-490.

[\[PubMed\]](#)

Effects of *Lactobacillus casei* Shirota, *Bifidobacterium breve* and oligofructose-enriched inulin on colonic nitrogen-protein metabolism in healthy humans. De Preter V, Vanhoutte T, Huys G *et al. American Journal of Physiology* **2007**;292: 358-368.

[\[Available on request\]](#)

Perioperative synbiotic treatment to prevent postoperative infectious complications in biliary cancer surgery: A randomized controlled trial. Sugawara G, Nagino M, Nishio H *et al. Annals of Surgery* **2006**;244(5):706-714.

[\[Free paper\]](#) [\[PubMed\]](#)

Randomized trial of dietary fibre and *Lactobacillus casei* administration for prevention of colorectal tumours. Ishikawa H, Akedo I, Otani T *et al. International Journal of Cancer* **2005**;116:762-767.

[\[Available on request\]](#)

Habitual intake of lactic acid bacteria and risk reduction of bladder cancer. Ohashi Y, Nakai S, Tsukamoto T *et al. Urologia Internationalis* **2002**;68:273-280.

[\[PubMed\]](#)

Preventive effect of a *Lactobacillus casei* preparation on the recurrence of superficial bladder cancer in a double blind trial. Aso Y, Akaza H, Kotake T *et al. European Urology* **1995**;27:104 – 109.

[\[PubMed\]](#)

Suppressing effect of *Lactobacillus casei* administration on the urinary mutagenicity arising from ingestion of fried ground beef in the human. Hayatsu H & Hayatsu T. *Cancer Letters* **1993**;73:173-179.

[\[PubMed\]](#)

8. Sistema immunitario

Diverse impact of a probiotic strain, *Lactocaseibacillus paracasei* Shirota, on peripheral mononuclear phagocytic cells in healthy Japanese office workers: a randomized, double-blind, controlled trial. Naito T, Morikawa M, Yamamoto-Fujimura M, Iwata A, Maki Ayaka, Kato-Nagaoka N, Oana K, Kiyoshima-Shibata J, Matsuura Y, Kaji R, Watanabe O, Shida K, Matsumoto S and Hori T. *Biosci Microbiota Food Health*. **2023**;42(1):65-72.

[\[Free paper\]](#) [\[Pubmed\]](#)

Effects of probiotic supplementation on natural killer cell function in healthy elderly individuals: a meta-analysis of randomized controlled trials. Gui Q, Wang A, Zhao X, *et al. Eur J Clin Nutr.* **2020**;1-8. doi:10.1038/s41430-020-0670-z

[\[Free paper\]](#) [\[PubMed\]](#)

Decreased number of days of fever detection and duration of fever with continuous intake of a fermented milk drink: a randomized, double-blind, placebo controlled study of elderly nursing home residents. Kushiro A, Shimizu K, Takada T, Kusunoki I and Aiba N. *Bioscience of Microbiota, Food and Health* **2019**;38(4), 151–157, 2019

[\[PubMed\]](#)

Effects of Short-Term Probiotic Ingestion on Immune Profiles and Microbial Translocation among HIV-1-Infected Vietnamese Children. Ishizaki A, Bi X, Nguyen LV, Matsuda K, Pham HV, Phan CTT, Khu DTK, Ichimura H. *Int J Mol Sci.* **2017** Oct 19;18(10):2185. doi: 10.3390/ijms18102185. PMID: 29048352; PMCID: PMC5666866.

[\[PubMed\]](#)

Immune response of healthy adults to the ingested probiotic *Lactobacillus casei* Shirota. Harbige LS, Pinto E, Allgrove J, Thomas LV. *Scandinavian Journal of Immunology* **2016**;84(6):353-364

[\[PubMed\]](#)

Effects of *Lactobacillus casei* Shirota ingestion on common cold infection and herpes virus antibodies in endurance athletes: a placebo-controlled, randomized trial. Gleeson M, Bishop N C, Struszcak L. *European Journal of Applied Physiology* **2016**;116(8):1555-63.

[\[Free paper\]](#) [\[PubMed\]](#)

Effect of Probiotic Supplement on Cytokine Levels in HIV-Infected Individuals: A Preliminary Study. Falasca K, Vecchiet J, Ucciferri C *et al.* *Nutrients* **2015**;28;7(10):8335-47

[\[Free paper\]](#) [\[PubMed\]](#)

Immunomodulatory effects of a probiotic drink containing *Lactobacillus casei* Shirota in healthy older volunteers. Dong H, Rowland I, Thomas LV, Yaqoob P. *European Journal of Nutrition* **2013**;52(8):1853-1863.

[\[PubMed\]](#)

Dysregulated circulating dendritic cell function in ulcerative colitis is partially restored by probiotic strain *Lactobacillus casei* Shirota. Mann ER, You J, Horneffer-van der Sluis V *et al* *Mediators Inflamm.* **2013**;2013:573576

[\[Free paper\]](#) [\[PubMed\]](#)

Daily *Lactobacillus casei* Shirota intake increases natural killer cell activity in smokers. Reale M, Boscolo P, Bellante V *et al.* *British Journal of Nutrition* **2011**;108(2):308-314.

[\[Available on request\]](#) [\[PubMed\]](#)

Flexible cytokine production by macrophages and T cells in response to probiotic bacteria. Shida K, Nanno M, Nagata S. *Gut Microbes* **2011**;2(2):109-114.

[\[PubMed\]](#)

Suppressive effect on activation of macrophages by *Lactobacillus casei* strain Shirota genes determining the synthesis of cell wall-associated polysaccharides. Yasuda E, Serata M, Sako T. *Applied & Environmental Microbiology* **2008**;74(15):4746-4755.

[\[Free paper\]](#)

Effects of a fermented milk drink containing *Lactobacillus casei* strain Shirota on the human NK-cell activity. Takeda K & Okumura K. *The Journal of Nutrition* **2007**;137(1): 791-793.

[\[Available on request\]](#) [\[PubMed\]](#)

Essential roles of monocytes in stimulating human peripheral blood mononuclear cells with *Lactobacillus casei* to produce cytokines and augment natural killer cell activity. Shida K, Suzuki T, Kiyoshima-Shibata J *et al.* *Clinical & Vaccine Immunology* **2006**;13(9):997-1003.

[\[Free paper\]](#) [\[PubMed\]](#)

Induction of interleukin-12 by *Lactobacillus* strains having a rigid cell wall resistant to intracellular digestion. Shida K, Kiyoshima-Shibata J, Nagaoka M, Watanabe K, Nanno M. *J Dairy Sci.* **2006** Sep;89(9):3306-17.

[\[Free paper\]](#) [\[PubMed\]](#)

Interleukin-12 is involved in the enhancement of human natural killer cell activity by *Lactobacillus casei* Shirota. Takeda K, Suzuki T, Shimada SI *et al. Clinical & Experimental Immunology* **2006**;146:109-115.

[\[PubMed\]](#)

Well-controlled proinflammatory cytokine responses of Peyer's patch cells to probiotic *Lactobacillus casei*. Chiba Y, Shida K, Nagata S *et al. Immunology* **2010**;130:352-362.

[\[PubMed\]](#)

Oral delivery of *Lactobacillus casei* Shirota modifies allergen-induced immune responses in allergic rhinitis. Ivory K, Chambers S, Pin C *et al. Clinical and Experimental Allergy* **2008**;38:1-8.

[\[Available on request\]](#) [\[PubMed\]](#)

Modulation of natural killer cell activity by supplementation of fermented milk containing *Lactobacillus casei* in smokers. Morimoto K, Takeshita T, Nanno M *et al. Preventive Medicine* **2005**;40:589-594.

[\[Available on request\]](#) [\[PubMed\]](#)

Effects of a fermented milk drink containing *Lactobacillus casei* strain Shirota on the immune system in healthy subjects. Nagao F, Nakayama M, Muto T, Okumura K. *Bioscience Biotechnology & Biochemistry* **2000**;64(12):2706-2708.

[\[Free paper\]](#)

9. Miscellanea (nutrizione enterale, sport, salute della cavità orale...)

Yearly changes in the composition of gut microbiota in the elderly, and the effect of lactobacilli intake on these changes. Amamoto R, Shimamoto K, Park S, Matsumoto H, Shimizu K, Katto M, Tsuji H, Matsubara S, Shephard RJ, Aoyagi Y. *Sci Rep.* **2021** Jun 17;11(1):12765.

[\[Free paper\]](#) [\[PubMed\]](#)

Very Long-Term Treatment with a *Lactobacillus* Probiotic Preparation, *Lactobacillus Casei* Strain Shirota, Suppresses Weight Loss in the Elderly. Ishikawa H, Mutoh M, Yoshimura K. *Nutrients* **2020**;12(6), 1599

[\[Free paper\]](#)

Effects of Daily Probiotics Supplementation on Anxiety Induced Physiological Parameters among Competitive Football Players. A.M.G.C.P. Adikari, Mahenderan Appukutty and Garry Kuan. *Nutrients* **2020**;12(7), 1920

[\[PubMed\]](#)

Effects of a probiotic drink containing *Lactobacillus casei* strain Shirota on dental plaque microbiota. Hu X, Huang Z, Zhang Y, Hong Y, Zheng Y. *J Int Med Res.* **2019**;47(7):3190-3202.

[\[PubMed\]](#)

Daily Intake of Fermented Milk Containing *Lactobacillus casei* Shirota (Lcs) Modulates Systemic and Upper Airways Immune/Inflammatory Responses in Marathon Runners.

Vaisberg M, Paixão V, Almeida EB, Santos JMB, Foster R, Rossi M, Pithon-Curi TC, Gorjão R, Momesso CM, Andrade MS, Araujo JR, Garcia MC, Cohen M, Perez EC, Santos-Dias A, Vieira RP, Bachi ALL. *Nutrients*. 2019 Jul; 11(7): 1678.

[\[PubMed\]](#)

The effect of probiotic *Lactobacillus casei* Shirota on knee osteoarthritis: a randomised double-blind, placebo-controlled clinical trial.

Lei M, Guo C, Wang D, Zhang C, Hua L. *Beneficial Microbes*. 2017 Oct 13;8(5):697-703.

[\[Abstract\]](#)

Effects of fermented soymilk with *Lactobacillus casei* Shirota on skin condition and the gut microbiota: a randomised clinical pilot trial.

Nagino T, Kaga C, Kano M *et al.* *Beneficial Microbes* 2017;9(2):209-218

[\[PubMed\]](#)

The effect of probiotic treatment on elderly patients with distal radius fracture: a prospective double-blind, placebo-controlled randomised clinical trial.

Lei M, Hua LM & Wang DW. *Beneficial Microbes* 2016;7(5):631-637

[\[PubMed\]](#)

Intake of a fermented soymilk beverage contains moderate levels of isoflavone aglycones enhances bioavailability of isoflavones in healthy premenopausal Japanese women: a double-blind, placebo-controlled, single-dose, crossover trial.

Nagino T, Kano M, Masuoka N *et al.* *Bioscience Microbiota, Food & Health* 2016;35(1):9-17

[\[Free paper\]](#) [\[PubMed\]](#)

The effect of five probiotic lactobacilli strains on the growth and biofilm formation of *Streptococcus mutans*.

Lin X, Chen X, Chen Y, Jiang W, Chen H. *Oral Diseases* 2015 Jan;21(1):e128-34. doi: 10.1111/odi.12257. Epub 2014 Jun 3. PMID: 24806217.

[\[Abstract\]](#)

The effect of a commercial probiotic drink containing *Lactobacillus casei* strain Shirota on oral health in healthy dentate people.

Sutula J, Coulthwaite LA, Thomas LV *et al.* *Microbial Ecology in Health & Disease* 2013;24:21003.

[\[PubMed\]](#)

***Lactobacillus casei* Shirota modulation of ammonia metabolism in physical exercise.**

Fuskevåg O-M, Annika Broster A, Davies N *et al.* *International Journal of Probiotics and Prebiotics* 2012;7(1):13-16.

[\[Abstract\]](#)

Effects of probiotic bacteria on *Candida* presence and IgA anti-*Candida* in the oral cavity of elderly.

Mendonça F, Santos S, Faria A *et al.* *Brazilian Dental Journal* 2012;23(5): 534-538

[\[ResearchGate\]](#)

Effects of synbiotic treatment on serum level of *p*-cresol in haemodialysis patients: a preliminary study. Nakabayashi I, Nakamura M, Kawakami K *et al* () *Nephrology Dialysis Transplantation* **2011**;26(3):1094-1098

[\[PubMed\]](#)

Probiotics affect the clinical inflammatory parameters of experimental gingivitis in humans. Slawik S, Staufenbiel I, Schilke R *et al*. *European Journal of Clinical Nutrition* **2011**;65(7):857-63

[\[PubMed\]](#)

Key role of teichoic acids on aflatoxin B1 binding by probiotic bacteria. Hernandez-Mendoza A, Guzman-de-Peña D, Garcia HS. *Journal of Applied Microbiology* **2009**;107(2):395-403.

[\[PubMed\]](#)

The influence of a probiotic milk drink on the development of gingivitis: a pilot study. Staab B, Eick S, Knofler G *et al*. *Journal of Clinical Periodontology* **2009**;36:850-856.

[\[PubMed\]](#)

Clinical safety of a *Lactobacillus casei* Shirota as a probiotic in critically ill children. Srinivasan R, Meyer R, Padmanabhan R, Britto J. *Journal of Pediatric Gastroenterology and Nutrition* **2006**;42:171-173.

[\[PubMed\]](#)

The *in vivo* use of stable isotope-labelled biomarkers lactose-[¹⁵N]ureide and [²H₄] tryosine to assess the effects of pro- and prebiotics on the intestinal flora of healthy human volunteers. De Preter V, Geboes K, Verbrugghe K *et al*. *British Journal of Nutrition* **2004**;92:439-446.

[\[PubMed\]](#)